

A NOTE ON SEMIGROUPS ON SPACES OF CONTINUOUS FUNCTIONS AND OF MEASURES

BÁLINT FARKAS

ABSTRACT. For Ω a Polish space we consider operator semigroups on the space $C_b(\Omega)$ of bounded, continuous functions and on the space $\mathcal{M}(\Omega)$ of bounded Borel measures. The important fact here is that the dual of $C_b(\Omega)$ with the strict topology is $\mathcal{M}(\Omega)$. We pursue the approach via bi-continuous semigroups, but we also show that the class of bi-continuous semigroups on $C_b(\Omega)$ with respect to the compact-open topology coincides with the class of equi-continuous semigroups with respect to the strict topology. If Ω is not Polish space this is not the case in general.

1. INTRODUCTION

The probably simplest example of a semigroup on the space $C_b(\mathbb{R})$, namely the shift semigroup, fails to be strongly continuous and even measurable with respect to the sup-norm. To overcome this and the probably the more important fact, namely the failure of strong continuity of many transition semigroups, several different methods have been developed, such as those of π -semigroups by Priola [14], weakly continuous semigroups by Cerrai [2]. One could even simply consider other locally convex topologies on $C_b(\mathbb{R})$ than the sup-norm-topology as, e.g., done by Dorroh, Neuberger [3]. A more abstract approach, we actually give preference to in this paper, is that of bi-continuous semigroups initiated by Kühnemund [6, 7].

Besides this progress, recently semigroups on spaces of measure have been attracting much attention, see Manca [13], Lant, Thieme [8]. It is not surprising however that such questions were addressed much earlier, for example by Sentilles [16], who studied operator semigroups of $C_b(\Omega)$ (bounded continuous functions) and on $\mathcal{M}(\Omega)$ (bounded measures), where Ω is a locally compact space. The important construction there was that of a *strict-topology*, which will be also crucial in this paper. In this respect we will rely on the paper Sentilles [15].

It is among the aims of the present note is to advertise bi-continuous semigroups that seem to be very much appropriate for dealing with one-parameter operator semigroups both on the space of bounded continuous functions or on the space of measures. One reason for giving distinction to this class of semigroups is that there is already a vast theory developed for them: there are generation, approximation and perturbation results (see Kühnemund [6], [7] and Farkas [10, 11]). More recently the Hille-Phillips functional calculus became also available in this setting, which can be used to prove convergence rates for rational approximation schemes and for efficient Laplace inversion formulas, see Jara [12]. We admit however that in many cases the class of bi-continuous semigroups is too small, and many interesting transition semigroups do not fall in this category. Regarding this, the notion of (weakly) stochastically continuous semigroups seems to be the most relevant, we refer the reader to Lant, Thieme [8]. There is moreover a recent paper by Kunze [4]

where integration of operator semigroups on spaces of bounded measurable functions, bounded continuous function or of continuous function vanishing at infinity is studied by means of duality pairs. This also allows the author to consider the Laplace transform and the adjoint of such semigroups in a fairly general setting.

Coming back to general bi-continuous semigroups, in this paper we show that under certain assumptions (which are satisfied for $C_b(\Omega)$, Ω Polish space) it is possible to consider bi-continuous semigroups, loosely speaking, on the dual $(X, \tau)'$, in particular the adjoint of a bi-continuous semigroup may be defined. This allows to consider bi-continuous semigroups on spaces of measures. We further show that these semigroups in many cases may be put in the theory of equicontinuous semigroups on locally convex spaces, but we also give an example, a rather pathological one, showing that this is not always possible. The results are taken from [9].

First we recall some terminology and set up the framework (see Kühnemund [6, 7]). For considering bi-continuous semigroups one needs first a Banach space $(X, \|\cdot\|)$ which is endowed with an additional locally convex topology τ . The two topologies need somehow be connected. We assume the following:

Hypothesis 1.1.

- (i) τ is Hausdorff and coarser than the norm-topology.
- (ii) The locally convex space (X, τ) is sequentially complete on τ -closed, norm-bounded sets.
- (iii) The dual space $(X, \tau)'$ is norming for $(X, \|\cdot\|)$, i.e.,

$$\|x\| = \sup_{\substack{\varphi \in (X, \tau)' \\ \|\varphi\| \leq 1}} |\varphi(x)|.$$

Now, τ -bi-continuous semigroups are defined as follows:

Definition 1.2. A semigroup $(T(t))_{t \geq 0}$ is called a *bi-continuous semigroup*, if

- (i) the orbit $t \mapsto T(t)x$ is τ -continuous for all $x \in X$,
- (ii) $t \mapsto T(t)$ is a norm-bounded function, say, on $[0, 1]$, in which case it is exponentially bounded.
- (iii) $(T(t))_{t \geq 0}$ is locally-bi-equicontinuous, which means that for a norm-bounded τ -null sequence x_n the convergence $T(t)x_n \rightarrow 0$ holds in the topology τ and uniformly in compact intervals.

The main features of this definition is that it first mixes properties of Banach spaces and a weaker notion of convergence, and second that it speaks about *convergent sequences*, which generally do not grasp the topological characteristics. As in (iii) above, for functions $T : \mathbb{R}_+ \rightarrow \mathcal{L}(X)$ we use the term “locally ...”, if the property “...” is satisfied for operators $T(t)$, t ranges over compact intervals of $\mathbb{R}_+$.

The illustrious examples of bi-continuous semigroups are those on the space $C_b(\Omega)$ when the continuity is required with respect to the compact-open topology τ_c . A more concrete instance is the shift semigroup on $C_b(\mathbb{R})$. To be more specific, consider Ω a locally compact, or a metrisable topological space, N.B. a completely regular space, (or even more generally a completely regular, k_f -space, i.e., a space Ω for which the continuity of a function $f : \Omega \rightarrow \mathbb{R}$ is decided already on compact sets). The linear space of continuous and bounded functions $f : \Omega \rightarrow \mathbb{R}$ is a Banach space if endowed with the supremum-norm $\|\cdot\|_\infty$. The additional topology that we consider is the *compact-open topology* τ_c generated by the following family of

semi-norms:

$$\left\{ p_K(f) := \sup_{x \in K} |f(x)| : K \subseteq \Omega \text{ compact} \right\}.$$

It is trivial that with $X = C_b(\Omega)$ and $\tau = \tau_c$ Hypothesis 1.1 is satisfied.

We show first how bi-continuous semigroups with respect to the compact open topology can be put in the framework of locally-equicontinuous semigroups, if Ω is a Polish space. For this purpose we introduce a further locally convex topology, called the *strict topology* (see Sentilles [15]). We give here three different constructions, which all coincide if Ω is a Polish space.

The first construction is a direct description of null-neighbourhoods. Consider the family $\mathcal{G}$ of all absolutely convex and absorbent sets $U \subseteq C_b(\Omega)$ which satisfy the following: for all $r > 0$ there exist a τ_c -neighbourhood V_r of 0, such that $U \cap B(0, r) \supseteq V_r \cap B(0, r)$. Then $\mathcal{G}$ is a base for a locally convex topology denoted by β_0 . Clearly, it is the finest locally convex topology which agrees with τ_c on all balls $B(0, r)$. It is called the *mixed topology*, see Wiweger [18].

For the other two constructions we need the Stone-Ćech compactification $\beta\Omega$ of the completely regular space Ω . The important fact is now the isometric isomorphism $C(\beta\Omega) \simeq C_b(\Omega)$, i.e., the fact that a bounded continuous function extends continuously from Ω to $\beta\Omega$ with the supremum being unchanged. For each compact set $K \subseteq \beta\Omega \setminus \Omega$ define

$$C_b^K(\Omega) := \{g : g \in C_b(\Omega), g|_K = 0\}.$$

Then consider the locally convex topologies β_K determined by the family of semi-norms

$$\mathcal{P}^K := \{\|f \cdot g\|_\infty : g \in C_b^K(\Omega)\}.$$

Now, β and β_1 are defined by inductive limits as follows

$$\beta := \varinjlim_{\substack{K \subseteq \beta\Omega \setminus \Omega \\ \text{compact}}} \beta_K \quad \text{and} \quad \beta_1 := \varinjlim_{\substack{K \subseteq \beta\Omega \setminus \Omega \\ \text{zero set}}} \beta_K.$$

For these topologies the following relations hold

$$\tau_c \leq \beta_0 \leq \beta \leq \beta_1 \leq \tau_n.$$

It is known that in all places “ $<$ ” is possible. However, when Ω is “nice”, some of the topologies may coincide. We have:

Theorem 1.3 (Sentilles [15]). *Let Ω either be a σ -compact, locally compact space, or a Polish space. Then*

- a) $\beta_0 = \beta = \beta_1$.
- b) $\beta = \mu(C_b(\Omega), \mathcal{M}(\Omega))$ the Mackey topology, where $\mathcal{M}(\Omega)$ denotes space of all bounded Baire-measures on Ω .
- c) A linear operator $T : C_b(\Omega) \rightarrow C_b(\Omega)$ is β -continuous, if and only if it is β -sequentially continuous. The same holds for linear functionals.
- d) Every $\varphi_n \in (C_b(\Omega), \beta)'$ weak*-null sequence is equicontinuous.

In general β and β_1 are referred to as the *sub-strict-topology* and the *strict topology*, respectively.

To close this section we present a further construction for mixing topologies. Let $\mathcal{P}$ denote a family of seminorms determining τ such that $p \leq \|\cdot\|$ for all $p \in \mathcal{P}$, and

$\|\cdot\| = \sup_{p \in \mathcal{P}} p(x)$. For $(p_n) \subseteq \mathcal{P}$ and $(a_n) \in c_0$, $a_n \geq 0$ consider the seminorm

$$\tilde{p}_{(p_n, a_n)}(x) = \sup_{n \in \mathbb{N}} a_n p_n(x).$$

The *mixed-topology* τ_m is determined by the family of seminorms

$$\tilde{\mathcal{P}} := \{\tilde{p}_{(p_n, a_n)} : p_n \in \mathcal{P}, a_n \in c_0\}.$$

It is clear, that

$$\tau \leq \tau_m \leq \tau_n,$$

where again τ_n denotes the norm-topology. A routine argument proves the following lemma:

Lemma 1.4. *A sequence $x_n \in X$ is convergent in the topology τ_m , if and only if it is norm-bounded and τ convergent.*

Proof. Note that it suffices to consider the case when the limit is 0. First let $x_n \xrightarrow{\tau_m} 0$ then clearly $x_n \xrightarrow{\tau} 0$. Suppose by contradiction that $\|x_n\|$ is unbounded. We may assume that $\|x_n\| \geq n$. For each x_n take $p_n \in \mathcal{P}$ such that $p_n(x_n) \geq \frac{n}{2}$. Thus for $\tilde{p}_{(p_k, 1/k)}$ we have

$$\tilde{p}_{(p_k, 1/k)}(x_n) = \sup_{\substack{k \in \mathbb{N} \\ k > 0}} \frac{1}{k} p_k(x_n) \geq \frac{1}{n} p_n(x_n) \geq \frac{1}{2},$$

which is a contradiction.

Conversely, assume that $x_n \xrightarrow{\tau} 0$ and x_n is norm-bounded. Take $a_n \in c_0$ and $p_n \in \mathcal{P}$. Then for $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that $|a_n| \leq \varepsilon$ whenever $n > N$. Thus

$$\tilde{p}_{(p_k, a_k)}(x_n) = \sup_{k \in \mathbb{N}} a_k p_k(x_n) \leq \sup_{0 \leq k \leq N} a_k p_k(x_n) + \sup_{k \geq N} a_k p_k(x_n) \leq C p'(x_n) + \varepsilon \|(a_k)\|,$$

which shows $x_n \xrightarrow{\tau_m} 0$. □

The following result translates the notion of bi-continuous semigroups to the language of mixed topologies.

Proposition 1.5. *The class of τ -bi-continuous semigroups and the class of τ_m -strongly continuous and locally sequentially τ_m -equicontinuous semigroups coincide.*

Proof. Let $(T(t))_{t \geq 0}$ be a τ -bi-continuous semigroup. Since $[0, 1] \ni h \rightarrow T(h)x$ is norm-bounded and τ -continuous for all x , we obtain by Lemma 1.4 that these orbits are also τ_m -continuous. The sequential τ_m -equicontinuity of the family

$$\{T(t) : t \in [0, t_0]\}$$

is simply a reformulation of Definition 1.2.(iii) in view of Lemma 1.4. For the converse, let $(T(t))_{t \geq 0}$ be a τ_m -strongly continuous and locally sequentially-equicontinuous semigroup. Then the orbits $[0, 1] \ni h \rightarrow T(h)x$ are norm-bounded since they are τ_m -continuous. The τ -strong continuity is immediate, while the local bi-equicontinuity is again trivial by Lemma 1.4. □

According to Wiweger [18], the topology τ_m coincides with the mixed topology under some abstract conditions, which are, in particular, satisfied for $(C_b(\Omega), \tau_c)$ and $(X', \sigma(X', X))$. Further he proves that in these cases τ_m is the finest locally convex topology which agrees with τ on norm-bounded sets, hence $\tau_m = \beta_0$.

2. DUALITY, MEASURES AND $C_b(\Omega)$

Given a τ -bi-continuous semigroup $(T(t))_{t \geq 0}$ on X , we would like to define the adjoint semigroup as a bi-continuous semigroup for the weak*-topology. However, for the validity of Hypothesis 1.1, we need to put further assumptions, which will be satisfied for $X = C_b(\Omega)$ and $\tau = \tau_c$. The established framework will be exploited in the last section.

Duality for (X, τ) . Let X° be the set of all norm-bounded linear functionals which are τ -sequentially continuous on norm-bounded sets of X . We endow this linear space with the operator norm inherited from the norm dual X' . We claim that it is indeed a Banach space, i.e., it is a closed linear subspace of X' . Take $\varphi_n \in X^\circ$ with $\|\varphi_n - \varphi\| \rightarrow 0$, where $\varphi \in X'$. We have to show that $\varphi \in X^\circ$. To this end, consider a norm-bounded τ -null sequence x_n . Then $\varphi(x_n) \rightarrow 0$ follows from

$$\begin{aligned} |\varphi(x_n)| &\leq |\varphi(x_n) - \varphi_k(x_n)| + |\varphi_k(x_n)| \leq \\ &\leq K\|\varphi - \varphi_k\| + |\varphi_k(x - x_n)| \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon, \end{aligned}$$

first by taking $k \in \mathbb{N}$ sufficiently large and then for fixed k using the continuity assumptions on φ_k .

Now, consider the Banach space X° equipped with the weak topology $\tau^\circ := \sigma(X^\circ, X)$. It is our aim to consider bi-continuous semigroups with respect to this topology on X° . For this purpose we have to verify the validity of Hypothesis 1.1. Clearly, τ° is Hausdorff since X separates the points of X° . For all $x \in X$ the operator norm of $x \in (X^\circ)'$ coincides with $\|x\|$, since X° is norming for X , and clearly $X = (X^\circ, \tau^\circ)'$. Hence

$$\|\varphi\| = \sup_{\substack{x \in X \\ \|x\| \leq 1}} |\varphi(x)|,$$

thus we see that $(X^\circ, \tau^\circ)'$ is norming for X° . Trivially, τ° is coarser than the norm-topology on X° . It remains to check only the τ° -sequential completeness on closed, norm-bounded sets, but this generally may fail to hold. At this point we have to put a condition on the space.

Hypothesis 2.1. Suppose that $X^\circ \cap \overline{B(0, 1)}$ is sequentially complete for $\sigma(X^\circ, X)$.

Below we prove that in the particular case $X = C_b(\Omega)$, which is our main concern now, we have $X^\circ = \mathcal{M}(\Omega)$, which is actually a well-known fact, and we basically recall a proof from Farkas [11]. Whereas we already point out here that to Alexandroff [1] it was known that $\mathcal{M}(\Omega)$ is $\sigma(\mathcal{M}(\Omega), C_b(\Omega))$ -sequentially complete if Ω is a Polish space, thus Hypothesis 2.1 is fulfilled in this situation.

The dual of $C_b(\Omega)$. Suppose that Ω is as in Theorem 1.3. Then by the considerations preceding Theorem 1.3 and by Theorem 1.3 itself it follows that $C_b(\Omega)^\circ$ coincides with the space of those functionals which are τ_c -continuous on norm-bounded sets. It remains therefore to determine these linear functionals, which can be easily achieved if Ω is a Polish space. The following considerations are taken almost verbatim from Farkas [11]. The dual of $C_b(\Omega)$ (as a Banach space) is isomorphic to the space $\mathcal{M}(\beta\Omega)$ of all bounded, Borel measures on the Stone-Ćech compactification $\beta\Omega$ of Ω , and the isomorphism is given by $\varphi(f) = \int_{\beta\Omega} f d\mu$. Since Ω is assumed to be topologically complete, it is known that Ω is G_δ set in $\beta\Omega$, see

Walker [17, Ch. 1]. Therefore it is possible to identify $\mathcal{M}(\Omega)$ with a subspace of $\mathcal{M}(\beta\Omega)$ in the following way:

$$\begin{aligned} \iota : \mathcal{M}(\Omega) &\rightarrow \mathcal{M}(\beta\Omega), \\ [\iota(\nu)](B) &:= \nu(\Omega \cap B) \quad \text{for all } \nu \in \mathcal{M}(\Omega) \text{ and } B \subseteq \beta\Omega \text{ Borel set.} \end{aligned}$$

Then ι is an injection with

$$\text{rg } \iota = \{\mu : \mu \in \mathcal{M}(\beta\Omega), |\mu|(\beta\Omega \setminus \Omega) = 0\}.$$

We show that a measure $\mu \in \mathcal{M}(\beta\Omega)$ gives rise to a linear functional $\varphi \in C_b(\Omega)'$ which is not only norm-continuous, but also τ_c -continuous on norm-bounded sets, if and only if it belongs to $\text{rg } \iota$, i.e. using the above identification, it is a regular Borel measure on Ω . Let $\mathcal{J}$ denote the set of these norm continuous-functionals, which is then a norm closed subspace of $\mathcal{M}(\beta\Omega)$. It is only to prove that $\varphi \in \mathcal{J}$ is associated with a measure in $\text{rg } \iota$, the other direction being easy. Assume the contrary, i.e., that $\varphi \in \mathcal{J}$ and $|\mu|(\beta\Omega \setminus \Omega) > c > 0$ for the corresponding μ . The measure $\mu|_\Omega$ is a Borel measure on the Polish space Ω , hence there is a compact set K_0 such that $|\mu|(\Omega \setminus K_0) < c/4$. Let $N \cup P = \beta\Omega$ be a Hahn-decomposition of μ on $\beta\Omega$. Then there are disjoint closed sets $F_1 \subseteq P \setminus \Omega$ and $F_2 \subseteq N \setminus \Omega$ with $\mu(P \setminus \Omega \setminus F_1) < c/8$ and $-\mu(N \setminus \Omega \setminus F_2) < c/8$. Take a further compact set $K \subseteq \Omega$ which is then closed in $\beta\Omega$. By Tietze's theorem we find a continuous function $f = f_{K \cup K_0, F_1, F_2}$ on $\beta\Omega$ with values in $[0, 1]$ satisfying $f(x) = 0$ for all $x \in K \cup K_0$ and $f(x) = (-1)^{i+1}$ when $x \in F_i$. We can write

$$\begin{aligned} \int_{\beta\Omega} f d\mu &= \int_{\Omega} f d\mu + \int_{\beta\Omega \setminus \Omega} f d\mu \\ &= \int_{K \cup K_0} f d\mu + \int_{\Omega \setminus (K \cup K_0)} f d\mu + \int_{P \setminus \Omega} f d\mu + \int_{N \setminus \Omega} f d\mu \\ &= \int_{K \cup K_0} f d\mu + \int_{\Omega \setminus (K \cup K_0)} f d\mu + \int_{F_1 \cup F_2} f d|\mu| + \int_{P \setminus \Omega \setminus F_1} f d\mu + \int_{N \setminus \Omega \setminus F_2} f d\mu \\ &\geq 0 - \frac{c}{4} + \frac{6c}{8} - \frac{c}{8} - \frac{c}{8} = \frac{c}{4} > 0. \end{aligned}$$

But this leads to contradiction by the arbitrariness of the compact set K and the τ_c -continuity of μ on the unit ball of $C_b(\Omega)$. We have therefore proved that $\mathcal{M}(\Omega)$ can be identified with $\mathcal{J}$. In combination with Theorem 1.3 this yields:

Theorem 2.2. *If Ω is a Polish space, then $(C_b(\Omega), \beta)' = C_b(\Omega)^\circ = \mathcal{M}(\Omega)$, and $\beta = \mu(C_b(\Omega), (C_b(\Omega), \beta)')$, i.e., $(C_b(\Omega), \beta)$ is a Mackey-space.*

Note that the above proof can be carried out for other classes of spaces Ω satisfying two requirements: (i) every finite Borel measure on Ω is automatically regular, (ii) Ω is Borel in $\beta\Omega$. For more general statement we refer the reader to Kunze [4].

3. SEMIGROUPS ON $C_b(\Omega)$ AND ON $\mathcal{M}(\Omega)$

First we start with an abstract construction for adjoint bi-continuous semigroups. Clearly a linear operator $B \in \mathcal{L}(X)$ which is also τ -sequentially continuous on norm-bounded sets admits an adjoint $B' \in \mathcal{L}(X')$ which leaves X° invariant. Take now a bi-continuous semigroup $(T(t))_{t \geq 0}$ on (X, τ) . Then the operators $T(t)^\circ$ form

a semigroup $(T^\circ(t))_{t \geq 0}$, which is τ° -strongly continuous by definition. The exponential boundedness of $(T^\circ(t))_{t \geq 0}$ is trivial. To establish the local bi-equicontinuity, we assume the following.

Hypothesis 3.1. Every $\varphi_n \in X^\circ$ norm-bounded τ° -null sequence is τ -sequentially equicontinuous on norm bounded sets.

To see the τ° -bi-equicontinuity of $(T^\circ(t))_{t \geq 0}$ under this hypothesis, take a norm-bounded τ° -null sequence and $x \in X$. For $t_0 > 0$ one has the τ -compactness of $\{T(t)x : t \in [0, t_0]\}$, thus by the equicontinuity of φ_n we have

$$[T^\circ(t)\varphi_n](x) = \varphi_n(T(t)x) \rightarrow 0$$

uniformly on $[0, t_0]$, which is the τ° -bi-equicontinuity of $(T^\circ(t))_{t \geq 0}$. Therefore if $(T(t))_{t \geq 0}$ is a bi-continuous semigroup.

Again the above hypothesis is satisfied for $C_b(\Omega)$ and τ_c . Indeed, if $\varphi_n \rightarrow 0$ in τ° , then the set $\{0, \varphi_n : n \in \mathbb{N}\}$ is weak*-compact hence is τ° -equicontinuous (see Theorem 2.2). These considerations show that it is always possible to take the dual $(T^\circ(t))_{t \geq 0}$ of a bi-continuous semigroup $(T(t))_{t \geq 0}$ on $C_b(\Omega)$. Instead of referring to Theorem 2.2, one could first use Prokhorov's Theorem to conclude that $\{0, \varphi_n : n \in \mathbb{N}\}$ is tight and then prove the following assertion, thus arriving to the equicontinuity of $\{\varphi_n : n \in \mathbb{N}\}$.

Proposition 3.2. *Let (Ω, d) be a Polish space. Then $\mathcal{K} \subseteq \mathcal{M}(\Omega)$ is tight and norm-bounded, if and only if it is τ_c -equicontinuous on norm bounded sets.*

Proof. Assume first the tightness and the norm-boundedness. Then the estimate

$$|\langle f, \varphi \rangle| \leq \int_K |f|d|\varphi| + \varepsilon \|f\|$$

for all $f \in C_b(\Omega)$ and $\varphi \in \mathcal{K}$ yields the equicontinuity of $\mathcal{K}$. For the contrary, let the family $\mathcal{K} \subseteq \mathcal{M}(\Omega)$ be τ_c -equicontinuous on norm-bounded sets. Assume by contradiction that there exists $\varepsilon > 0$ such that for all $K \subseteq \Omega$ compact there exist a functional $\varphi_K \in \mathcal{K}$ with $|\varphi_K|(\Omega \setminus K) \geq \varepsilon$. For $K \subseteq \Omega$ compact, the sets $K_n := \{x : d(x, K) < 1/n\}$ are open, and one has $\bigcap_{n=1}^\infty K_n = K$. Take $n \in \mathbb{N}$ such that $|\varphi_K|(\Omega \setminus K_n) \geq \varepsilon/2$. Define the function $f|_K := 0$ and $f|_{\Omega \setminus K_n} := 1$ and extend it to the whole of Ω with values in $[0, 1]$ (Tietze's theorem). Then

$$\int_\Omega f d|\varphi_K| \geq \int_{\Omega \setminus K_n} f d|\varphi_K| \geq \varepsilon.$$

This contradicts the equicontinuity assumption since in all τ_c -neighbourhood of 0 one finds such f . $\square$

Remark. a) Note that for a Polish space Ω , since $(C_b(\Omega), \beta)$ is a Mackey-space by Theorem 2.2, the β -equicontinuous sets of $\mathcal{M}(\Omega) = (C_b(\Omega), \beta)'$ are exactly the relative weak*-compact sets. So first by first showing the above (trivial) assertion, Theorem 3.2, one obtains the proof of Prokhorov's theorem.

b) From the above it follows that if a bounded operator T is τ_c -(sequentially)-continuous on sup-norm bounded sets, then its adjoint T' leaves $X^\circ = \mathcal{M}(\Omega)$ invariant.

Lemma 3.3. *Let $T : \mathbb{R}_+ \rightarrow \mathcal{L}(C_b(\Omega))$ be a τ_c -strongly continuous function and $\mathcal{K} \subseteq \mathcal{M}(\Omega)$ a weak*-compact set. Then the map*

$$[0, +\infty) \times \mathcal{K} \ni (t, \nu) \mapsto T'(t)\nu \in \mathcal{M}(\beta\Omega)$$

is continuous if we take the weak-topology $\sigma(\mathcal{M}(\beta\Omega), C_b(\Omega))$ on $\mathcal{K}$ and on $\mathcal{M}(\beta\Omega)$.*

Proof. Let $t \in [0, +\infty)$ and $\nu \in \mathcal{K}$ fixed. Then

$$\begin{aligned} |\langle f, T'(t)\nu - T'(t')\nu' \rangle| &= |\langle T(t)f, \nu \rangle - \langle T(t')f, \nu' \rangle| \leq \\ &|\langle T(t)f, \nu - \nu' \rangle| + |\langle T(t)f - T(t')f, \nu' \rangle| \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \quad \text{for all } f \in C_b(\Omega). \end{aligned}$$

The last inequality follows by the continuity assumptions and by Prokhorov's theorem for $t' \in U$ and $\nu' \in V$ with appropriate neighbourhoods U, V of t and ν . $\square$

A straightforward consequence of this lemma and the fact that the set $[0, t_0] \times \mathcal{K}$ is compact for the product topology is the following:

Lemma 3.4. *Let $T : [0, +\infty) \rightarrow \mathcal{L}(C_b(\Omega))$ be a τ_c -strongly continuous function. For a weak*-compact set $\mathcal{K} \subseteq \mathcal{M}(\Omega)$ and $t_0 > 0$ the set of measures*

$$\{T'(t)\nu : t \in [0, t_0], \nu \in \mathcal{K}\}$$

is $\sigma(\mathcal{M}(\beta\Omega), C_b(\Omega))$ -compact.

Lemma 3.5. *Let $T : \mathbb{R}_+ \rightarrow \mathcal{L}(C_b(\Omega))$ be a τ_c -strongly continuous semigroup consisting of operators that are τ_c -continuous on norm-bounded sets. For a norm-bounded, weak*-compact set $\mathcal{K} \subseteq \mathcal{M}(\Omega)$ and $t_0 > 0$ the set of measures*

$$\{T'(t)\nu : t \in [0, t_0], \nu \in \mathcal{K}\}$$

is tight.

Proof. The assertion follows immediately from Lemma 3.4, taking into account the arguments preceding Lemma 3.3 concerning the invariance of $\mathcal{M}(\Omega)$. $\square$

Proposition 3.6. *Let $T : \mathbb{R}_+ \rightarrow \mathcal{L}(C_b(\Omega))$ be τ_c -strongly continuous and locally norm-bounded. Suppose that $T(t)$ takes norm-bounded τ_c -null sequences into τ_c -null sequences. Then for all compact set $K \subseteq \Omega$ and $\varepsilon > 0$, there exists $M > 0$ and $K' \subseteq \Omega$ compact such that*

$$\sup_{x \in K} |(T(t)f)(x)| \leq M \sup_{x \in K'} |f(x)| + \varepsilon \|f\|_\infty \quad .$$

holds uniformly for t in compact intervals in $\mathbb{R}$. In particular, it is locally- β -equicontinuous, or, which the same, it is τ_c -bi-equicontinuous.

Proof. Let $\varepsilon > 0$, $t_0 > 0$ and $K \subseteq \Omega$ be a compact set. Take a compact set $K' \subseteq \Omega$ such that $|T'\delta_x|(\Omega \setminus K') \leq \varepsilon$ for all $t \in [0, t_0]$ and $x \in K$. Such compact set exists by Lemma 3.5. We then obtain

$$\begin{aligned} \sup_{x \in K} |T(t)f(x)| &= \sup_{x \in K} \left| \int_{\Omega} f dT'(t)\delta_x \right| \leq \sup_{x \in K} \int_{K'} |f| d|T'(t)\delta_x| + \sup_{x \in K} \int_{\Omega \setminus K'} |f| d|T'(t)\delta_x| \leq \\ &\leq \sup_{t \in [0, t_0]} \|T(t)\| \cdot \sup_{x \in K'} |f(x)| + \varepsilon \|f\|, \end{aligned}$$

which is the assertion. $\square$

A variant of this result has been obtained independently by Markus Kunze, see ???. Let us summarise the above in the following two assertions.

Theorem 3.7. *Let Ω be a Polish space and consider the set $\mathcal{F}$ of τ_c -strongly continuous semigroups $(T(t))_{t \geq 0}$ on $C_b(\Omega)$ for which each $T(t)$ is τ_c -sequentially continuous on sup-norm bounded sets. Then the class of τ_c -bi continuous semigroups and the class of β -locally equicontinuous, β -strongly continuous semigroups both coincide with $\mathcal{F}$.*

Theorem 3.8. *Let Ω be a Polish space and $(T(t))_{t \geq 0}$ a τ_c -bi-continuous semigroup on $C_b(\Omega)$. Then the semigroup $(T(t)^\circ)_{t \geq 0}$ defines as $T(t)^\circ := T(t)'|_{\mathcal{M}(\Omega)}$ is a bi-continuous semigroup on the space of bounded, regular Borel measures $\mathcal{M}(\Omega)$ with respect to the topology $\sigma(\mathcal{M}(\Omega), C_b(\Omega))$.*

The crucial ingredient, i.e., that a bounded, linear operator leaves $\mathcal{M}(\Omega)$ invariant if and only if it is τ_c continuous on norm-bounded sets, or which is the same, if it is β_0 -continuous, has long since been known, and has found several independent proofs, see Sentilles [15], Priola [14], Farkas [11] or Manca [13]. A proof in a more general case is given in Kunze [4].

A surprising fact is that though τ_c is generally not metrisable, the continuity of norm-bounded linear operators on norm-bounded sets can be described by convergent sequences. It is clear that “completeness” or “compactness” play an important role here, whereas some kind of countability has been also overall assumed (metric or σ -compact spaces). Indeed, a simplest non-countable space gives rise to a counterexample to Theorem 3.7, when Ω is not a Polish space. More specifically we construct below a bi-continuous semigroup which is not β_0 -locally-equicontinuous.

Example 3.9. Let $\Omega = \omega_1$ the first uncountable ordinal number and v be the order topology. Since any $\alpha \in \omega_1$ has cofinality less than ω_0 , we see that Ω is first countable and normal. Suppose that $f_n \rightarrow 0$ in the topology τ_c . We claim that there exists $\alpha \in \omega_1$ such that $f_n \rightarrow 0$ uniformly on $[\alpha, \omega_1)$. Suppose the contrary, i.e., for all $\alpha < \omega_1$ there exists $k \in \mathbb{N}$, $k > 0$ such that for all $N \in \mathbb{N}$ there exists $n \geq N$ and $x \in [\alpha, \omega_1)$ with $|f_n(x)| > 1/k$. For all $\alpha \in \omega_1$ we have $k_\alpha \in \mathbb{N}$ and we may assume that $k_{\alpha_\xi} = k$ for a cofinal sequence $\alpha_\xi \in \omega_1$. By induction we choose a sequence

$$x_{\alpha_{\xi_1}} < x_{\alpha_{\xi_1}} + 1 < x_{\alpha_{\xi_2}} < x_{\alpha_{\xi_2}} + 1 < \dots < x_{\alpha_{\xi_j}} < x_{\alpha_{\xi_j}} + 1 < \dots$$

with $f_{n_j}(x_{\alpha_{\xi_j}}) > 1/k$. Since $K := \{\lim_{j \rightarrow \infty} x_{\alpha_{\xi_j}}, x_{\alpha_{\xi_j}} : j \in \mathbb{N}\}$ is compact and

$$\sup_{y \in K} |f_{n_j}(y)| \geq \frac{1}{k} \quad \text{for all } j \in \mathbb{N},$$

we arrived to a contradiction. Thus we have the existence of $\alpha \in \omega_1$ as asserted above. Now, consider the family $\{[\xi, \omega_1) : \xi > \alpha\}$, which has the finite intersection property and thus by compactness possesses an accumulation point $x \in \beta\Omega$. All f_n extends to the Stone-Ćech compactification $\beta\Omega$ and $|f_n(y)| < \varepsilon$ for all $y \in [\alpha, \omega_1)$ if $n \geq N$. Take $n \in \mathbb{N}$, $n \geq N$. By the continuity of f_n on $\beta\Omega$ we have a neighbourhood U of x such that for all $y \in U$

$$|f_n(y) - f_n(x)| \leq \varepsilon.$$

There exist $\xi \in (\alpha, \omega_1)$ with $\emptyset \neq U \cap [\xi, \omega_1) \ni z$, so

$$|f_n(x)| \leq |f_n(x) - f_n(z)| + |f_n(z)| \leq \varepsilon + \varepsilon.$$

Thus $f_n(x) \rightarrow 0$, which shows that δ_x is τ_c -sequentially-continuous (on norm-bounded sets). However, it is clear that this is not τ_c -continuous on norm-bounded sets. Consider now the C_0 -semigroup $(T(t))_{t \geq 0}$ generated by the bounded operator $A := \mathbf{1} \otimes \delta_x$. Since A is idempotent the semigroup T takes the form

$$T(t) = I - A + e^t A.$$

This semigroup is bi-continuous but none of $T(t)$, $t > 0$ is τ_c continuous on norm bounded sets.

REFERENCES

- [1] A. D. Alexandroff, *Additive set-functions in abstract spaces*, Mat. Sb. **13** (**55**) (1943), 169–238.
- [2] S. Cerrai, *A Hille-Yosida theorem for weakly continuous semigroups*, Semigroup Forum **49** (1994), 349–367.
- [3] J. R. Dorroh, J. W. Neuberger, *Lie generators for semigroups of transformations on a Polish space*, Electron. J. Differential Equations **1** (1993), 1–7.
- [4] M. Kunze, *Very weak integration of transition semigroups*, Ulmer Seminare **12** (2007), 285–299.
- [5] M. Kunze, personal communication, April 2008.
- [6] F. Kühnemund: *Bi-Continuous Semigroups on Spaces with Two Topologies: Theory and Applications*, Ph.D. thesis, Universität Tübingen, 2001.
- [7] F. Kühnemund: *A Hille-Yosida Theorem for bi-continuous semigroups*, Semigroup Forum **67** (2003), 205–225.
- [8] T. Lant, H. R. Thieme, *Markov transition functions and semigroups of measures*, Semigroup Forum **74** (2007), no. 3, 337–369.
- [9] B. Farkas, *Perturbations of bi-continuous semigroups*, Ph.D. Thesis, Eötvös Loránd University, 2003.
- [10] B. Farkas, *Perturbations for bi-continuous semigroups*, Studia Math. **161**(2) (2004), 147–161.
- [11] B. Farkas *Perturbations of bi-continuous semigroups on $C_b(H)$ with applications to the Ornstein-Uhlenbeck semigroup*, Semigroup Forum **68** (2004), 87–107.
- [12] P. Jara, *Rational approximation schemes for bi-continuous semigroups*, J. Math. Anal. Appl., to appear; doi:10.1016/j.jmaa.2008.02.068
- [13] L. Manca, *Kolmogorov equations for measures*, preprint. <http://arxiv.org/abs/math/0703654>
- [14] E. Priola, *On a class of Markov type semigroups in spaces of uniformly continuous and bounded functions*, Studia Math. **136** (1999), 271–295.
- [15] F. D. Sentiilles, *Bounded continuous functions on a completely regular space*, Trans. Amer. Math. Soc. **168** (1972), 311–336.
- [16] F. D. Sentiilles, *Semigroups of operators in $C(S)$* , Canad. J. Math. **22** (1970), 47–154.
- [17] R. C. Walker, *The Stone-Čech Compactification*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 83, Springer-Verlag, New York, 1974.
- [18] A. Wiweger, *Linear spaces with mixed topology*, Studia Math. **20** (1961), 47–68.

TECHNISCHE UNIVERSITÄT DARMSTADT, FACHBEREICH MATHEMATIK
 SCHLOSSGARTENSTR. 7, D-64289 DARMSTADT (GERMANY)
 E-mail address: `farkas@mathematik.tu-darmstadt.de`